

Tear Film and Ocular Surface Surfactants

Lars Bräuer and Friedrich P. Paulsen*

Department of Anatomy and Cell Biology, Martin Luther University of Halle-Wittenberg, Halle/Saale, Germany

Abstract: The hydrophobic surfactant proteins B (SP-B) and C (SP-C) are tightly bound to phospholipids. These proteins play important roles in maintaining the surface tension-lowering properties of pulmonary surfactant. Surfactant protein A (SP-A) and D (SP-D) are extremely hydrophilic and are thought to have a role in recycling surfactant and especially in improving host defense in the lung. Moreover, SP-A supports the hydrophobic surfactant proteins during surfactant sub-type assembly and inhibits secretion of lamellar bodies into the alveolar space. During recent years surfactant proteins have also been detected at locations outside the lung such as the lacrimal apparatus. In this review, the latest information regarding SP function and regulation in the human lacrimal system, the tear film and the ocular surface is summarized with regard to mucous epithelial integrity, rheological and antimicrobial properties of the tear film, tear outflow, certain disease states and possible therapeutic perspectives.

Keywords: Surfactant proteins, ocular surface, tear film, lacrimal apparatus.

INTRODUCTION

Each year worldwide 55 million ocular injuries occur (WHO-Program for the prevention of blindness). Ocular trauma and corneal ulcerations due to infection are the most significant causes of corneal blindness yearly leading to 1.6 million new instances of bilateral blindness, 2.3 million instances of serious limitation in visual acuity, and 19 million instances of unilateral blindness. In Germany, approximately 30 million people wear glasses or contact lenses. In the USA, the number approaches 120 million. In 1995, the number of radial keratotomies (RK) reached 250,000 in the USA (National Eye Institute). Meanwhile, RK has been almost completely superseded by LASIK (laser assisted in situ keratomileusis) (1.55 million operations in the USA in the year 2000). These numbers have increased appreciably over the last seven years. The laser market has a yearly growth rate of 10-25%. Although LASIK is a great deal safer and associated with fewer complications than RK and photoreactive keratoectomy, standardized numbers on the occurrence of LASIK related complications have not yet become available. To date, the only relevant publications have been based on small case numbers at individual centers.

Due to the difficulties involved in treating corneal blindness once it has developed, it is, from a medical point of view, well worth the effort to search for therapeutic strategies that could improve healing early and effectively, reduce infections and diminish corneal scarring in order to reduce the number of instances of persistent corneal blindness.

In ophthalmological practice, bacterial keratitis and conjunctivitis are among the most frequently seen problems. They represent an important component of numerous infections of the eye, particularly in patients after penetrating

corneal injury, after lengthy periods of contact lens wear, after refractive corneal surgery or under immune suppression therapy [1-4]. The most frequent causal agents are the gram positive pathogen *Staphylococcus aureus* and the gram negative bacteria *Pseudomonas aeruginosa* [5, 6], but also viruses, e.g. the *Herpes simplex* virus, can instigate keratitis. Bacterial by-products and toxins as well as the host's inflammatory reaction often bring about extensive tissue damage with persistent scarring or even loss of sight [1].

It has long been known that tear fluid contains various anti-bacterially active substances. These substances are synthesized by various cells found in the lacrimal glands, the accessory lacrimal glands, the conjunctiva and the cornea. Of these antimicrobial substances, lysozyme [7], mainly effective against gram positive bacteria, has been most extensively described and studied. Further substances include the enzyme β -Lysin, which destroys bacterial cell membranes working synergistically with lysozyme [8], the iron-binding protein Lactoferrin, which inhibits growth of *Bacillus subtilis*, *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Pseudomonas aeruginosa* [9, 10], as well as complement factors of the alternate pathway, which are activated directly by microbial products, e.g. endotoxins [11]. In the past few years, the research group surrounding Bernhard Redl has been able to better characterize the protein lipocalin present in the lacrimal fluid and has recently demonstrated that tear-specific lipocalin is bacteriostatic to *Escherichia coli* and various fungi *via* iron deprivation [12]. Recent research has further shown that lacrimal fluid contains secretory phospholipase A₂, an enzyme which, along with lysozyme and β -lysin is particularly effective against gram-positive bacteria [13, 14].

A further mechanism of defence is represented by antimicrobial peptides. It is assumed that some of these peptides are able to kill microorganisms directly *via* formation of pores. For the remainder of these peptides the mechanism of activity has not yet been fully elucidated. The best-characterized group of antimicrobial peptides are the human

*Address correspondence to this author at the Department of Anatomy and Cell Biology, Martin Luther University Halle-Wittenberg, Faculty of Medicine, Grosse Steinstr. 52, D-06097 Halle (Saale), Germany; Tel: +49 345 5571707; Fax: +49 345 5571700; E-mail: friedrich.paulsen@medizin.uni-halle.de

defensins that can be subdivided into two groups, α -defensins and β -defensins. Further antimicrobial peptides that have been described on the ocular surface include BPI (bactericidal permeability-increasing protein), CAP37 (heparin-binding protein), LL37 (human cationic antimicrobial protein/hCAP-18), LEAP1 (liver expressed antimicrobial peptide 1 or heptacidin) and LEAP 2 and angiogenin. A current review of antimicrobial peptides at the ocular surface has been presented by McDermott [15].

SURFACTANT PROTEINS

The surfactant-associated proteins A and D (SP-A and SP-D) have been extensively described in research related to the lung. Immunological functions in both the non-specific and specific immune defence systems have been ascribed to them. SP-A and SP-D are representatives of the collectin-family of the C-type, among with numerous other molecules of known immunological function may be counted.

According to the current understanding of the mechanism of C-type collectins, exposed microbiological carbohydrates (on the surface of diverse microorganisms) bind to a carbohydrate recognition domain inherent to the proteins. In this manner, opsonisation and acceleration of the defence reaction to the microorganism is achieved.

SP-A is a 28-36 kDa soluble protein first recognized in type II pneumocytes of the pulmonary epithelium and is expressed by these cells. SP-A isolated from bronchoalveolar lavage specimens is mainly present in the form of surfactant-lipid-aggregates. Furthermore, the presence of SP-A augments the reduction in surface tension induced by SP-B in pulmonary alveoli. Nonetheless, deletion of the SP-A gene in mice does not lead to a decline in lung stability, albeit the protein inhibition of surfactant reveals an important functional relevance of SP-A [16]. SP-A bind to type II pneumocytes and to immune cells as well as to some extent to macrophages [17]. Beyond this, it has been demonstrated that SP-A deficient mice exhibit a weakened defence to various pulmonary pathogens [18].

SP-D is a 43 kDa protein synthesized not only by type II pneumocyte epithelium, but also by various cells of the respiratory tract and mucous cells of the stomach. It is characterized by a very high structural similarity and, correspondingly, homology to SP-A and other mammalian specific lectins [19]. The lectin domain of SP-D, mediated by calcium ions, bind to carbohydrates and lipids, playing a role in the innate immune defence particularly against bacterial, viral and fungiform pathogens. It interacts directly with a number of microorganisms such as influenza virus A [20], *Pseudomonas aeruginosa* and *Escherichia coli* [21, 22]. In addition, its important role in lipid homeostasis of the lung could be demonstrated through the selective deletion of SP-D in mice [23]. SP-D deficient mice develop alveolar lipidosis, which is accompanied by the activation of macrophages and an increased activation of metalloproteinases [24]. Beyond this, the SP-D deficient mice develop emphysema-like pathological changes reflecting the critical role of the surfactant proteins in the regulation of pulmonary inflammation.

The first description of proteins in organic extracts of extracellular surfactant and lamellar bodies was presented by Phizackerley *et al.* in 1979 [25]. The characterization and purification of these proteins proved quite difficult due to

their high hydrophobicity and low molecular weights. Finally, it could be demonstrated that pulmonary surfactant contains at least two different small molecular weight hydrophobic proteins, known as SP-B and SP-C [26, 27]. Using non-reducing preparation-conditions the molecular weights of SP-B by means of SDS-PAGE have been estimated at approximately 15-18 kDa, whereas under reducing conditions a suitably separable product of approximately 7 kDa was obtained [28, 29]. SP-B, post-translationally produced from a preform is also able to form oligomeres of various sizes *via* disulfide bridges [29]. SP-C, utilising 33-35 amino acids, is one of the smallest and at the same time most hydrophobic proteins known, with a molecular weight of 4-6 kDa [30, 31]. Its primary translation product consists of 191 amino acids. As does SP-B, SP-C also undergoes extensive post-translational modification, for example, glycosylation, acylation or esterification with fatty acids [32-35]. In contrast to the sugar binding collectin-like SP-A and SP-D, SP-B and SP-C are decisive in the formation and stability of surface active layers (membranes) and beyond this are a prerequisite for the absorption of phospholipids at the air-fluid boundary [36, 37]. Due to their hydrophobic properties, SP-B and SP-C are of great pathophysiological relevance in acute respiratory distress syndrome (ARDS) as diverse plasma proteins inhibit the formation of pulmonary surfactants and thus decrease alveolar surface tension [38]. In this context it has been reported that the deletion of SP-B in newborn rabbits leads to serious disturbance of the affected surfactant and injury to the alveolar surface of therewith engendered respiratory stress in the rabbit [39]. The function of SP-C is comparable to SP-B in that it is characterized by its direct influence and interaction with biological and phase interfaces; it namely reduces surface tension and increases the ability to absorb the surfactant, like an anchor mediating between the phospholipid layer and the aqueous phase.

SURFACTANT PROTEINS OF THE TEAR FILM, OCULAR SURFACE AND LACRIMAL APPARATUS

The Meibomian glands of the eyelids are responsible for the production of the superficial lipid components of the tear film which counteract evaporative tendencies of the deeper tear film layers directly adjacent to and protecting the ocular surface [40]. The lacrimal glands, along with the accessory glands of the lids, express the watery components of the tear film. These play a role not only in moistening of the ocular surface, but also in defending against potentially pathogenic microorganisms [13]. The mucous components of the tear film consist for the most part of secretory mucins produced in conjunctival goblet cells, the lacrimal and accessory lacrimal glands, as well as of membrane bound mucins synthesized by conjunctival and corneal epithelial cells [41-45]. Membrane bound mucins may be split from the epithelial surface *via* a process known as shedding, and thus may also enter the aqueous component of the tear film [44, 45].

For a long time it has been supposed that superficially active substances, similar to the surfactant system of the lung, are of importance not only in tear film but also in the auditory tube and on the skin [46]. Various studies have documented the presence of surfactant-associated proteins A, B and D in the auditory tube [47] and of SP-A, B, C and D in the skin [48]. The presence of SP-D has already been described in tear film and lacrimal glands [49-51]. Ni *et al.*

[52] was able to show that beyond this, SP-D is present in the cornea of mice and has protective effects against keratitis caused by *P. aeruginosa*. Although Dobbie *et al.* [53] had already presented immunohistochemical evidence of the presence of SP-A in the human lacrimal gland, these findings were pushed into the background by Stahlmann *et al.* [50] and Ni *et al.* [52] as the latter authors could not demonstrate SP-A in tear fluid of mice and their findings were taken to be conclusive for humans as well. Our group has now been able to show that not only SP-D, but also the surfactant-associated protein A, along with SP-B and SP-C is present at the ocular surface, in the lacrimal apparatus and in tears [54, 55]. Thereby, all of the surfactant-associated proteins (SP's) proved to be detectable at the mRNA and protein levels throughout the lacrimal apparatus and on the ocular surface in all tissues studied (lacrimal glands, conjunctiva, cornea, lacrimal ducts) as well as in tear fluid itself. Interestingly, SP-A and SP-C in tear fluid, and SP-C in all examined tissues, show an expression pattern differing from that of lung surfactant proteins. This is probably due to tissue specific post-translational or post-transcriptional modifications of the proteins and may lead to differences in the spectrum of activity of the surfactant proteins. SP-A, -B, -C and -D are found in the acinar cells of the lacrimal gland and the accessory tear ducts, in the conjunctival epithelial cells and in columnar epithelial cells (particularly apically) as well as in serous portions of seromucous glands in the tear ducts and the accessory tear ducts. Goblet cells do not produce any of the four SP's. In contrast to tear fluid, the aqueous humor does not show the presence of SP's under physiological conditions [54]. In healthy corneal specimens, the two collectins SP-A and SP-D are only found as a thin film in the apical region of the superficial cells. SP-B and SP-C are immunohistochemically undetectable within and on the surface of the cornea [55]. In contrast, pathologically altered corneae show a completely different distribution pattern. In association with herpes keratitis and corneal ulceration (caused by *S. aureus*), SP-A and SP-D are found in the vicinity of the lesion sites as well as in invading immune cells; in the corneae of patients with keratoconus the entire epithelium and the endothelium react positively. Cultured corneal and conjunctival epithelial cell lines also produce SP-A and SP-D [54]. Stimulation of corneal and conjunctival epithelial cell lines [56, 57] with various cytokines and bacterial supernatants lead to an increase of both collectins. Thus, in addition to SP-D which has already been reported to be present in tear film [49, 50, 52], there are also SP-A, SP-B and SP-C proteins in the lacrimal fluid and on the ocular surface.

CONCLUSIONS

Based on the present status of knowledge including that gleaned from animal experimental studies on SP-D [52, 58] it can be assumed that the collectins, SP-A and SP-D, are involved in pathological processes at the ocular surface and function here, as well as in the efferent tear ducts, in the service of non-specific natural immune defence and in the activation of the adaptive immune system. As a substance intrinsic to tears they protect the ocular surface in conjunction with immunoglobulin A (IgA), defensins and mucins against infection by *P. aeruginosa*, *S. aureus* and other pathogenic microbes [15, 52, 59-62]. In the efferent tear ducts they are also active in conjunction with IgA, antimicro-

bial peptides and mucins in preventing the formation of dacryocystitis [54, 55, 63-70].

In view of the fact that the hydrophobic surfactant proteins SP-B and SP-C have an expanding influence on the surface tension of the air-liquid interface on top of the alveolar lining layer, a similar effect could be discussed in relation to the tear film and the tear fluid at the human ocular surface and efferent tear ducts. In this context we provide a completed and modified scheme of the tear film which now contains the evidenced and investigated surfactant proteins with respect to their physicochemical properties and discussed putative functions in tears, the lacrimal apparatus and at the ocular surface (Fig. 1). This view demonstrates the small hydrophobic surfactant proteins B and C embedded into the lipid component of the tear film, orientated regarding to their amphiphilic character. Furthermore, the water-soluble and polymerizable collectin-like surfactant proteins A and D are arranged within the aqueous component of the tear film along with the already known different secreted and shedded mucins. This hypothetical model supports possible functions of surfactant proteins with regard to severe pathologies of the ocular surface e.g. dry eye syndrome and bacterial or viral infections. Dysfunctions of the complex ocular surfactant system would probably lead to considerable visual impairment and disturbances of the immune defence system of the ocular surface and lacrimal apparatus. Hypothetically, absence of the small hydrophobic surfactant proteins B and C could result in alterations of tear film stability and as a consequence in interruption of the tear film itself leading to symptoms of dry eye syndrome such as for example dry spots. Disturbed production or lack of the immunologically important surfactant proteins A and D could lead to impairment of innate and adaptive immunity at the ocular surface and in the lacrimal system resulting in various bacterial and/or viral infections.

PERSPECTIVES

By means of the development of a gene construct for all four surfactant proteins, which includes open reading frames for mature and preforms of the proteins, a prerequisite has been formed for the establishment of a system of expression *via* which all four SP's can be manufactured using recombinant methods. The thus manufactured recombinant SP's could serve in functional studies on questions pertaining to the ocular surface and the lacrimal system. Some recombinant SP's have already been manufactured (for a review see [71, 72]). With the software programs and methods available today, it is possible to create a reliable model of the 3D-structure of proteins and to identify the active sites by using comparative protein modelling or threading. Blocking or modification of putative active sites could alter the functionality of the proteins and lead to completely new proteins at a functional level. Up to now there is no information on the ocular surface and lacrimal system of surfactant-deficient mice, which are already available for all four SP's [23, 73-75]. In studies on conditions of chronic infection of the ocular surface such as that found, for example, in dry eyes, it would be of interest to know whether individual SP formation is increased or decreased. An increase in surfactant formation over a lengthy period of time would very possibly have negative consequences for the body as a whole. In this context it has been demonstrated that SP-D exhibits proat-




Fig. (1). The figure demonstrates a modified and completed scheme of the present understanding of the tear film in accordance to that recently published by Paulsen (2006). It displays the arrangement of the water-soluble and putative immunologically important surfactant proteins A and D within the aqueous phase of the tear film. Furthermore, the small hydrophobic surfactant proteins B and C are illustrated. They are embedded into the lipid-phase of the tear film with respect to their physicochemical properties. Moreover, most of the already known and still investigated mucins of the ocular surface and tear film are shown along with SP-A and SP-D.

herogenous activity in mice [76]. Studies on the surfactant-protein system of the ocular surface have just begun and lead us to expect interesting new discoveries in regard to future therapeutic perspectives.

ACKNOWLEDGEMENT

This work was supported by grants from the Deutsche Forschungsgemeinschaft (DFG) – program grants PA 738/9-1, PA 738/9-2 and PA 738/1-5, the BMBF – Wilhelm Roux Program, Halle, Germany – program grants FKZ 09/17, as well as Sicca Forschungsförderung of the Association of German Ophthalmologists

LIST OF ABBREVIATIONS

SP	= surfactant protein
WHO	= World Health Organization
RK	= radial keratotomy
LASIK	= laser assisted in situ keratomileusis
CAP37	= heparin-binding protein
LL37	= human cationic antimicrobial protein
hCAP18	= human cationic antimicrobial protein
LEAP1	= liver expressed antimicrobial peptide 1/heptacidin
LEAP2	= liver expressed antimicrobial peptide 2

SDS-PAGE = sodium dodecyl sulfate-polyacrylamide gel electrophoresis

ARDS = acute respiratory distress syndrome

mRNA = messenger ribonucleic acid

IgA = immunoglobulin A

REFERENCES

- [1] Baum J. Infections of the eye. *Clin Infect Dis* 1995; 21: 479-88.
- [2] Gritz DC, Whitcher JP. Topical issues in the treatment of bacterial keratitis. *Int Ophthalmol Clin* 1998; 38: 107-14.
- [3] Brennan NA, Chantal Coles ML. Extended wear in perspective. *Invest Ophthalmol Vis Sci* 1997; 74: 609-23.
- [4] Levartovsky S, Rosenwasser G, Goodman D. Bacterial keratitis following laser in situ keratomileusis. *Ophthalmology* 2001; 108: 321-5.
- [5] Fleiszig SMJ, Efron N, Pier GB. Extended contact lens wear enhances *Pseudomonas aeruginosa* adherence to human corneal epithelium. *Invest Ophthalmol Vis Sci* 1992; 33: 2908-16.
- [6] O'Callaghan RJ, Cellegan MC, Moreau JM, *et al.* Specific roles of alpha-toxin and beta-toxin during *Staphylococcus aureus* corneal infection. *Infect Immun* 1997; 65: 1571-8.
- [7] Fleming A. On a remarkable bacteriolytic element found in tissues and secretions. *Proc R Soc B* 1922; 93: 306-17.
- [8] Pleyer U, Baatz H. Antibacterial protection of the ocular surface. *Ophthalmologica* 1997; 211 (Suppl 1): 2-8.
- [9] Oram J, Reiter B. Inhibition of bacteria by lactoferrin and other iron-chelating agents. *Biochim Biophys Acta* 1979; 170: 351-3.
- [10] Kijlstra A. The role of lactoferrin in the nonspecific immune response on the ocular surface. *Regul Immunol* 1990; 3: 193-7.
- [11] Fearson D, Austen K. The alternative pathway of complement: A system for host resistance to microbial infection. *N Engl J Med* 1980; 303: 259-61.

- [12] Fluckinger M, Haas H, Merschak P, Glasgow BJ, Redl B. Human tear lipocalin exhibits antimicrobial activity by scavenging microbial siderophores. *Antimicrob Agents Chem* 2004; 48: 3367-72.
- [13] Qu X-D, Lehrer RI. Secretory phospholipase A₂ is the principal bactericide for staphylococci and other gram-positive bacteria in human tears. *Infect Immun* 1998; 66: 2791-9.
- [14] Girgis DO, Dajcs JJ, O'Callaghan RJ. Phospholipase A₂ activity in normal and *Staphylococcus aureus*-infected rabbit eyes. *Invest Ophthalmol Vis Sci* 2003; 44: 197-202.
- [15] McDermott AM. Defensins and other antimicrobial peptides at the ocular surface. *Ocul Surf* 2004; 2: 229-47.
- [16] Korfhagen TR, Bruno MD, Ross GF, *et al.* Altered surfactant function and structure in SP-A gene targeted mice. *Proc Natl Acad Sci USA* 1996; 93: 9594-9.
- [17] van Iwaarden F, Welmers B, Verhoef J, Haagsman HP, van Golde LM. Pulmonary surfactant protein A enhances the host-defense mechanism of rat alveolar macrophages. *Am J Respir Cell Mol Biol* 1990; 2: 91-8.
- [18] Levine AM, Kurak KE, Bruno MD, Stark JM, Whitsett JA, Korfhagen TR. Surfactant protein-A-deficient mice are susceptible to *Pseudomonas aeruginosa* infection. *Am J Respir Cell Mol Biol* 1998; 19: 700-8.
- [19] Crouch E, Wright JR. Surfactant proteins a and d and pulmonary host defense. *Annu Rev Physiol* 2001; 63: 521-54.
- [20] Hartshorn KL, Crouch E, White MR, *et al.* Pulmonary surfactant proteins A and D enhance neutrophil uptake of bacteria. *Am J Physiol* 1998; 274: L958-69.
- [21] Sastry K, Ezekowitz RA. Collectins: pattern recognition molecules involved in first line host defense. *Curr Opin Immunol* 1993; 5: 59-66.
- [22] Ferguson JS, Voelker DR, McCormack FX, Schlesinger LS. Surfactant protein D binds to *Mycobacterium tuberculosis* bacilli and lipoarabinomannan *via* carbohydrate-lectin interactions resulting in reduced phagocytosis of the bacteria by macrophages. *J Immunol* 1999; 163: 312-21.
- [23] Botas C, Poulain F, Akiyama J, *et al.* Altered surfactant homeostasis and alveolar type II cell morphology in mice lacking surfactant protein D. *Proc Natl Acad Sci USA* 1998; 95: 11869-74.
- [24] Wert SE, Yoshida M, LeVine AM, *et al.* Increased metalloproteinase activity, oxidant production, and emphysema in surfactant protein D gene-inactivated mice. *Proc Natl Acad Sci USA* 2000; 23: 5972-7.
- [25] Phizackerley PJ, Town MH, Newman GE. Hydrophobic proteins of lamellated osmiophilic bodies isolated from pig lung. *Biochem J* 1979; 183: 731-6.
- [26] Yu SH, Chung W, Olafson RW, Harding PG, Possmayer F. Characterization of the small hydrophobic proteins associated with pulmonary surfactant. *Biochim Biophys Acta* 1987; 921: 437-48.
- [27] Whitsett JA, Hull WM, Ohning B, Ross G, Weaver TE. Immunologic identification of a pulmonary surfactant-associated protein of molecular weight = 6000 daltons. *Pediatr Res* 1986; 20: 744-49.
- [28] Yu SH, Wallace D, Bhavnani B, Enhorning G, Harding PG, Possmayer F. Effect of reconstituted pulmonary surfactant containing the 6000-dalton hydrophobic protein on lung compliance of prematurely delivered rabbit fetuses. *Pediatr Res* 1988; 23: 23-30.
- [29] Whitsett JA, Ohning BL, Ross G, *et al.* Hydrophobic surfactant-associated protein in whole lung surfactant and its importance for biophysical activity in lung surfactant extracts used for replacement therapy. *Pediatr Res* 1986; 20: 460-7.
- [30] Possmayer F. A proposed nomenclature for pulmonary surfactant-associated proteins. *Am Rev Respir Dis* 1988; 138: 990-8.
- [31] Hawgood S, Shiffer K. Structures and properties of the surfactant-associated proteins. *Annu Rev Physiol* 1991; 53: 375-394.
- [32] Voorhout WF, Veenendaal T, Haagsman HP, *et al.* Intracellular processing of pulmonary surfactant protein B in an endosomal/lysosomal compartment. *Am J Physiol* 1992; 263: 479-86.
- [33] Glasser SW, Korfhagen TR, Perme CM, Pilot-Matias TJ, Kister SE, Whitsett JA. Two SP-C genes encoding human pulmonary surfactant proteolipid. *J Biol Chem* 1988; 263: 10326-31.
- [34] Jacobs KA, Phelps DS, Steinbrink R, *et al.* Isolation of cDNA clone encoding a high molecular weight precursor to a 6-kDa pulmonary surfactant-associated protein. *J Biol Chem* 1987; 262: 9808-11.
- [35] Curstedt T, Johansson J, Barros-Soederling J, *et al.* Low-molecular-mass surfactant protein type I. The primary structure of a hydrophobic 8-kDa polypeptide with eight half-cystine residues. *Eur J Biochem* 1988; 172: 521-5.
- [36] Yu SH, Possmayer F. Role of bovine pulmonary surfactant-associated proteins in the surface-active property of phospholipid mixtures. *Biochim Biophys Acta* 1990; 1046: 233-41.
- [37] Notter RH, Shapiro DL, Ohning B, Whitsett JA. Biophysical activity of synthetic phospholipids combined with purified lung surfactant 6000 dalton apoprotein. *Chem Phys Lipids* 1987; 44: 1-17.
- [38] Spragg, RG, Gilliard, N, Richman, P. In: Robertson B, van Golde LMG Batenburg JJ, Eds. *Pulmonary Surfactant: From molecular biology to clinical practice.* Amsterdam, Elsevier. 1992; 685-703.
- [39] Kobayashi T, Nitta K, Takahashi R, Kurashima K, Robertson B, Suzuki Y. Activity of pulmonary surfactant after blocking the associated proteins SP-A and SP-B. *J Appl Physiol* 1991; 71: 530-6.
- [40] McCulley JP, Shine WE. Meibomian gland function and the tear lipid layer. *Ocul Surf* 2003; 1: 97-106.
- [41] Gipson IK, Hori Y, Argueso P, *et al.* Character of ocular surface mucins and their alteration in dry eye disease. *Ocul Surf* 2004; 2: 131-48.
- [42] Jumblatt MM, McKenzie RW, Steele PS, Emberts CG, Jumblatt JE. MUC7 expression in the human lacrimal gland and conjunctiva. *Cornea* 2003; 22: 41-5.
- [43] Paulsen F, Langer G, Hoffmann W, Berry M. Human lacrimal gland mucins. *Cell Tissue Res* 2004; 316: 167-77.
- [44] Paulsen F. Cell and molecular biology of human lacrimal gland and nasolacrimal duct mucins. *Int Rev Cytol* 2006; 249: 229-79.
- [45] Paulsen F, Berry M. Mucins and TFF peptides of the tear film and lacrimal apparatus. *Prog Histochem Cytochem* 2006; 41: 1-53.
- [46] Ridsdale RA, Palaniyar N, Possmayer F, Harauz G. Formation of folds and vesicles by dipalmitoylphosphatidylcholine monolayers spread in excess. *J Membr Biol* 2001; 180: 21-32.
- [47] Paananen R, Sormunen R, Glumoff V, van Eijk M, Hallman M. Surfactant proteins A and D in Eustachian tube epithelium. *Am J Physiol Lung Cell Mol Physiol* 2001; 281: 660-7.
- [48] Mo YK, Kankavi O, Masci PP, *et al.* Surfactant protein expression in human skin: evidence and implications. *J Invest Dermatol* 2007; 127: 381-6.
- [49] Madsen J, Kliem A, Tornøe I, Skjodt K, Koch C, Holmskov U. Localization of lung surfactant protein D on mucosal surfaces in human tissues. *J Immunol* 2000; 164: 5866-70.
- [50] Stahlman MT, Gray ME, Hull WM, Whitsett JA. Immunolocalization of surfactant protein-D (SP-D) in human fetal, newborn, and adult tissues. *J Histochem Cytochem* 2002; 50: 651-60.
- [51] Akiyama J, Hoffman A, Brown C, *et al.* Tissue distribution of surfactant proteins A and D in the mouse. *J Histochem Cytochem* 2002; 50: 993-6.
- [52] Ni M, Evans DJ, Hawgood S, Anders EM, Sack RA, Fleiszig SM. Surfactant protein D is present in human tear fluid and the cornea and inhibits epithelial cell invasion by *Pseudomonas aeruginosa*. *Infect Immun* 2005; 73: 2147-56.
- [53] Dobbie JW, Tasiaux N, Meijers P, *et al.* Lamellar bodies in synovocytes, mesothelium and specific epithelia as possible site of auto-antigen in rheumatoid disease. *Br J Rheumatol* 1994; 33: 508-19.
- [54] Bräuer L, Kindler C, Jäger K, *et al.* Detection of surfactant proteins A and D in human tear fluid and the human lacrimal system. *Invest Ophthalmol Vis Sci* 2007; 48: 3945-53.
- [55] Bräuer L, Bürgermann J, Johl M, Pleyer U, Tsokos M, Paulsen F. Detection and localization of the hydrophobic surfactant proteins B and C in human tear fluid and the human lacrimal system. *Curr Eye Res* 2007; 32: 931-8.
- [56] Araki-Sasaki K, Ohashi Y, Sasabe T, *et al.* An SV40-immortalized human corneal epithelial cell line and its characterization. *Invest Ophthalmol Vis Sci* 1995; 36: 614-21.
- [57] Diebold Y, Calonge M, Enriquez de Salamanca A, *et al.* Characterization of a spontaneously immortalized cell line (IOBA-NHC) from normal human conjunctiva. *Invest Ophthalmol Vis Sci* 2003; 44: 4263-74.
- [58] McCormick CC, Hobden JA, Balzli CL, *et al.* Surfactant protein D in *Pseudomonas aeruginosa* keratitis. *Ocul Immunol Inflamm* 2007; 15: 371-9.
- [59] Masinick SA, Montgomery CP, Montgomery PC, Hazlett LD. Secretory IgA inhibits *Pseudomonas aeruginosa* binding to cornea and protects against keratitis. *Invest Ophthalmol Vis Sci* 1997; 38: 910-8.

- [60] McNamara NA, Van R, Tuchin OS, Fleiszig SM. Ocular surface Epithelia expression mRNA for human beta defensin-2. *Exp Eye Res* 1999; 69: 483-90.
- [61] McNamara NA, Andika R, Kwong M, Sack RA, Fleiszig SM. Interaction of *Pseudomonas aeruginosa* with human tear fluid components. *Curr Eye Res* 2005; 30: 517-25.
- [62] Kwong MSF, Evans DJ, Ni M, Cowell BA, Fleiszig SMJ. Human tear fluid protects against *Pseudomonas aeruginosa* keratitis in a murine experimental model. *Infect Immun* 2007; 75: 2325-32.
- [63] Paulsen F, Thale A, Kohla G, *et al.* Functional anatomy of human duct epithelium. *Anat Embryol* 1998; 198: 1-12.
- [64] Paulsen F, Pufe T, Schaudig U, *et al.* Detection of natural peptide antibiotics in human nasolacrimal ducts. *Invest Ophthalmol Vis Sci* 2001; 42: 2157-63.
- [65] Paulsen F, Pufe T, Schaudig U, *et al.* Protection of human efferent tear ducts by antimicrobial peptides. *Adv Exp Med Biol* 2002; 506: 547-53.
- [66] Paulsen F, Corfield A, Hinz M, *et al.* Characterization of mucins in human lacrimal sac and nasolacrimal duct. *Invest Ophthalmol Vis Sci* 2003; 44: 1807-13.
- [67] Paulsen F, Corfield A, Hinz M, *et al.* Tränenabfluss – Bedeutung von Muzinen und TFF-Peptiden. *Ophthalmologe* 2004; 101: 19-24.
- [68] Paulsen F, Varoga D, Steven P, Pufe T. In: Zierhut M, Stern ME, Sullivan DA, Eds. *Immunology of Lacrimal Gland and Tear Film*. London, Taylor & Francis. 2005; 97-104.
- [69] Paulsen F. In: Weber R, Keerl R, Schaefer SD, Della Rocca RC, Eds. *Atlas of Lacrimal Surgery*. Berlin, Heidelberg, New York, Springer. 2007; 1-13.
- [70] Paulsen F. Functional anatomy and immunological interactions of ocular surface and adnexa. *Dev Ophthalmol* 2008; 41: 21-35.
- [71] Pfister RH, Soll RF. New synthetic surfactants: the next generation? *Biol Neonate* 2005; 87: 338-44.
- [72] Sorensen GL, Husby S, Holmskov U. Surfactant protein A and surfactant protein D variation in pulmonary disease. *Immunobiology* 2007; 212: 381-416.
- [73] Akinbi HT, Breslin JS, Ikegami M, *et al.* Rescue of SP-B knockout mice with a truncated SP-B proprotein. *J Biol Chem* 1997; 272: 9640-7.
- [74] Li G, Siddiqui J, Hendry M, *et al.* Surfactant protein-A-deficient mice display an exaggerated early inflammatory response to a β -resistant strain of influenza A virus. *Am J Respir Cell Mol Biol* 2002; 26: 277-82.
- [75] Glasser SW, Burhans MS, Korfhagen TR, *et al.* Altered stability of pulmonary surfactant in SP-C-deficient mice. *Proc Natl Acad Sci* 2001; 98: 6366-71.
- [76] Sorensen GL, Madsen J, Kejling K, *et al.* Surfactant protein D is proatherogenic in mice. *Am J Physiol Hear Circ Physiol* 2006; 290: H2286-94.

Received: June 06, 2008

Revised: July 29, 2008

Accepted: July 31, 2008

© Bräuer and Paulsen; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.